The irritable baby

Neil J. Davies

- p0330 The constantly irritable infant is often referred to by primary health care providers in a somewhat offhanded way as being colicky, as if that somehow represents the final word on the subject. 'Infantile colic' is a term which is used by various health care professions to describe the persistent, often violent crying which sometimes characterizes an otherwise healthy and thriving baby. As many as 1 in 5 parents report a problem with infant irritability or crying in the first 3 months of life but less than 1 in 20 with problem crying have an organic cause (Hiscock & Jordan 2004).
- Although there have been a number of theories proposed p0335 as to the etiology of colic, the literature remains fraught with difficulties in definition, methodological problems and numerous claims as to both etiology and management that are anecdotal (Hewson et al 1987). Various medical authors believe colicky behavior to be a reaction pain, probably of intestinal origin (Behrman et al 2007, Hull & Johnston 1987, Illingworth 1985). That this characteristic irritable behavior attributed to colic which is uncomplicated by other disease is a result of pain is also supported by researchers within both chiropractic and medicine (Biedermann 1992, Gutmann 1987, Kloughart et al 1989, Nilsson 1985). While the exact mechanisms for this pain remain to some extent obscure, the work of Sato (1980) and Budgell (1998) has shed considerable light on the subject, suggesting a neurophysiological mechanism that supports the clinical observations of Kloughart et al (1989) in their multicentered research trial which demonstrated a striking success rate for resolution of colicky symptoms in young infants after chiropractic management.
- p0340 Infantile colic probably remains best conceptualized as the end result of a complex transaction between the infant and their environment with multiple factors potentially responsible for the crying and distress (Hewson et al 1987). However, for a subgroup of infants, early excessive crying may evolve into a more generalized 'persistent mother-infant distress' syndrome (Leung & Lemay 2004). Taken together, these factors open the way for wide-ranging management options for the chiropractor.

Uncomplicated colic

The purpose of using the term 'uncomplicated colic' is to clearly p0345 signify in the clinical record that there are no factors such as protein intolerance, carbohydrate intolerance, or obvious parenting errors that are complicating the colicky presentation. These complications are discussed later in this chapter.

Clinical presentation

The symptoms that are characteristic of the typical colicky p0350 infant begin abruptly: the cry is loud and more or less continuous, persisting for several hours at the same time each day, usually late in the afternoon or early in the evening; the face sometimes becomes flushed or circumoral pallor develops; the abdomen may be distended and tense with the legs drawn up; the feet are often found to be cold and the hands bunched into tightly held fists. The paroxysm may simply spontaneously terminate or relief is sometimes obvious after the passage of flatus or fecus (Barr 1998, Behrman et al 2007).

Diagnostic evaluation

Reaching a diagnostic conclusion of uncomplicated colic is a p0355 process of elimination. The child with colicky symptoms must be carefully evaluated for any clinical signs that may suggest underlying disease. The parents/caregivers must also be carefully questioned in order to establish that there are no parenting errors, environmental factors, feeding difficulties or family history that may be contributing to the child's irritable disposition. Only after all these factors have been taken into consideration and ruled out, can the chiropractor safely assume that the child indeed has a diagnosis of uncomplicated colic.

Table 5.1 lists some of the more common clinical conditions p0360 which may complicate the colicky presentation and their symptomatic characteristics.

s0010

s0015

s0020



The irritable baby

t0010 Table 5.1 Clinical indicators that may imply underlying disease in irritable infant presentations

Body system	Symptom/sign	Associated condition
Integument	Maculopapular rash on trunk, neck, face and upper arms Excoriated erythematous skin on the buttocks and anogenital region Pustules over face and trunk with general lymphadenopathy Erythematous rash which desquamates and forms pustules in an otherwise healthy term infant	Cow's milk or soy protein intolerance Congenitally low pH of urine and other body fluids Congenital infection Congenital candidiasis
Gastrointestinal	 Bloating, frequent passage of flatus, constipation and/or diarrhea Sudden onset of paroxysms of severe colicky pain in a previously unaffected infant which get steadily worse over a few hours Profuse, explosive diarrhea which has a sweet smell and tests positive for disaccharides Projectile vomiting immediately following feeding preceded by 'golf ball' reverse peristaltic waves seen under tangential lighting of the upper abdomen Constant small volume vomiting which has a sour smell Unchanged nappies (diapers) Swallowed air Overfeeding, underfeeding, or inappropriately clothed for the prevailing temperature 	Cow's milk or soy protein intolerance Intussusception or other bowel obstruction Lactose intolerance Pyloric stenosis Gastroesophageal reflux Parenting error Nursing technique Parenting error
Respiratory	Crackles/wet sounds without obvious dyspnea Crackles with fever and dyspnea	Cow's milk or soy protein intolerance Infection
Genitourinary	Fever with decreased fluids out	Renal infection
Head and neck	Hair in the child's eye Fever Nasal congestion, postnasal drip and reddened tonsils Head held constantly to one side	Corneal irritation Otitis Chronic catarrh Congenital sternomastoid tumor

Chiropractic management s0025

- p0365 Clinical studies with pediatric patients presenting with visceral conditions in general have shown somewhat variable responses to chiropractic management (Ali et al 2002, Bronfort et al 2002, Fallon & Edelman 1998, Froehle 1996, Gemmell & Jacobsen 1989, Lebouef et al 1991, Nilsson & Christiansen 1988, Reed et al 1994). In terms of infantile colic, a number of studies have reported that chiropractic is effective in producing symptomatic relief in a relatively short timeframe (Davies & Jamison 2007, Klougart et al 1989, Mercer & Nook 1999, Wiberg et al 1999). However, one controlled clinical trial conducted in Norway concluded that 'chiropractic spinal manipulation' was no more effective than placebo in the treatment of infantile colic (Olafsdottir et al 2001). The growing weight of evidence in the scientific literature suggests that chiropractic management is safe and certainly warranted in cases where the irritability of babies can be classified as 'uncomplicated colic.'
- p0370 Once the diagnosis has been established to a clinical certainty, there are two major aspects involved in chiropractic management. The first is to assess and precisely correct the patterns of neurological dysafferentation. In years past the focus of the chiropractic profession has been largely upon the upper cervical complex and the T8-T12 region as representing the most likely anatomical location of the subluxation in colicky babies (Anrig & Plaugher 1998, Davies 2000). Since the advent of a more neurological approach to chiropractic, it now needs to be said that the anatomical location is less predictable than

we first thought and driven more by patterns of cortical dysafferentation and the associated dural tension and cerebrospinal fluid flow dynamics rather than nerve root factors.

The second aspect of care is to counsel the parents and p0375 caregivers carefully in relation to feeding practices, positive reinforcement, and the use of contingent music therapy. The correction of neurological dysafferentation, while having been shown to be effective in resolving the distressing symptoms associated with uncomplicated colicky presentations (Meadow & Smithells 1975), is not the totality of patient management as the important factors referred to above cause frustration for both the infant and mother, creating a positive feedback cycle which in turn perpetuates irritability and recurrence of the subluxation complex, presumably via the viscerosomatic, somatosomatic, and psychosomatic reflex pathways. Provision of adequate, practical counseling to the nursing mother is an essential element in the chiropractic management program (Taubman 1990).

Use of contingent music and differential reinforcement

s0030

It has been shown by clinical research that there is a significant p0380 behavioral aspect to children with uncomplicated colic presentations that can be considerably varied by the use of contingent music and differential reinforcement (Larson & Allyon 1990). Parents with colicky children should be carefully counseled to turn on music when the infant is quiet and alert for a period

The irritable baby

CHAPTER 5

exceeding 30 seconds. In addition, at such times, reinforcement should be given in the form of parental attention, including making eye contact, talking softly, rocking, walking, playing with the baby and generally being loving and affectionate. When crying resumes, parents should turn off music and attend to the infant's needs. This behavioral approach, which is designed to compete with crying by reinforcing quiet alertness with music and parental attention, has been demonstrated to reduce crying time by as much as 75% (Larson & Allyon 1990). This technique of behavioral modification therapy is simple to employ within the home environment and is a valuable adjunct to chiropractic management.

Protein intolerance and allergy

p0385 This syndrome, despite extensive research reported in the scientific literature, continues to be underdiagnosed, frequently misdiagnosed, and even more frequently mismanaged. All too often cow's milk intolerance is misdiagnosed as uncomplicated colic and subsequent management programs fail accordingly. In chiropractic, such a mistake in diagnosis leads to overadjusting with all the attendant problems that accompany that form of mismanagement. Physicians and chiropractors alike continue to tell mothers that breastfeeding protects their child from cow's milk allergy and intolerance despite the wealth of studies which have been available in the scientific literature for over a decade demonstrating both the presence and clinical effect of cow's milk protein in human breastmilk (e.g. Barau & Dupont 1994, Chandra 1997, deJong et al 1998, Goldborough & Francis 1983, Gustafsson et al 1992, Host et al 1988, McCarty & Frick 1983, Wilson et al 1990).

s0040 Pathophysiology

- p0390 The physiology of the human gut in the first year of life varies significantly from that of the older child in that the permeability is much greater. By the end of the first year of life the permeability becomes rapidly more adult-like. Because of this increased permeability, during the first year of life, proteinaceous substances can make their way across the gut wall into the bloodstream where they can produce an IgE-mediated antigen-antibody response (Plebani et al 1990, Wilson & Hamburger 1988). There is no single substance in cow's milk that produces the allergenic reaction. Casein, alpha-lactalbumin and beta-lactalbumin all show a high proportion of positive reactions (Savilahti & Kuitunen 1992) with specific IgE antibodies to these substances having been recovered from infants allergic to cow's milk (Bjorksten et al 1983). It has been further demonstrated that children with cow's milk intolerance have an even greater permeability of the small intestine than is the case with non-atopic children of the same age (Schrander et al 1992). Some cases of cow's milk intolerance, however, cannot be shown to be IgE mediated and therefore it seems apparent that the child's symptoms arise from a non-immune source (Foucard 1985).
- p0395 While intolerance to cow's milk is typically seen in babies where there is a positive family history of atopic disease (Foucard 1985, Wilson & Hamburger 1988), it is not necessarily

always the case. Certainly, a negative family history does not automatically eliminate a diagnosis of cow's milk intolerance.

Clinical presentation

The presentation of cow's milk intolerance shows a consistent p0400 pattern of symptoms of which those related to the gastrointestinal system predominate. Allergic reactions involving the skin, respiratory system, and neurological system are also common (Burr et al 1993; Businco et al 1985; Chandra & Hamed 1991; Chandra et al 1989; Iacono et al 1998; Kahn et al 1988, 1989; Lucas et al 1990; Nadasi 1992).

Diagnosis

In order to make a presumptive clinical diagnosis of cows milk p0405 allergy, a 'triad' of symptoms, which must include the gastrointestinal, respiratory, and integumentary systems, must be present. The majority of children who end up with a confirmed diagnosis of cow's milk allergy will present initially with this typical 'triad' of symptoms, namely gastrointestinal disturbance, skin rash and respiratory 'wet sounds'. While a small number present atypically, it is probably safe to say that all children with cow's milk allergy will at least have gastrointestinal symptoms and one or more of the other symptoms shown in Table 5.2.

Once the possibility of cow's milk allergy is considered, it is p0410 wise to arrange appropriate allergy tests and an evaluation of total serum IgE. When found in combination, a positive radioallergosorbent test (RAST) to cow's milk and an elevated total serum IgE is as reliable as laboratory investigations get in assisting the clinician to confirm a diagnosis of cow's milk allergy. Certainly, a combination of typical symptomatic pattern and positive laboratory results will confirm the diagnosis in a good percentage of cases.

 Table 5.2 Common symptoms associated with cow's milk allergy
 t0015

 and intolerance in infants

Body system	Symptom/sign
Gastrointestinal	Bloating, frequent passage of flatus and intractable crying/distress with pulling up of the legs Chronic diarrhea, constipation or an alternating pattern of both
Integument	Maculopapular rash which may occur anywhere on the body, but is found most commonly on the face, neck, trunk, buttocks and upper arms. Sometimes it is only seen in body fossae and within the enclosed areas of skin folds Eczema
Respiratory	Crackles/wet sounds without obvious dyspnea Wheezing and rhinitis
Neurological	Disturbed sleep pattern with frequent waking, crying at night, short sleep cycles and long non-rapid eye movement stage 1 (NREM1) sleep

61

s0045

s0050



p0415 It should be borne in mind when performing laboratory tests, however, that a second type of reaction to cow's milk which is really a non-immune intolerance will return negative laboratory results while the symptomatic picture remains strongly evidential of the diagnosis. Under these circumstances, a clinical diagnosis should be made and the patient managed accordingly. A successful clinical outcome is all the diagnostic confirmation that is needed.

The irritable baby

s0055 Management

- p0420 Chiropractic management of the infant who reacts to cow's milk protein is the same for IgE-mediated and non-immune intolerances. In both cases, diagnostic confirmation is dependent upon clinical outcome. The management of the breastfed infant is simple. The mother is counseled to avoid all food substances containing dairy products and to ensure a fluid intake of approximately 1.5-2 liters per day for the trial period of 15 days. The baby should be evaluated for evidence of neurological dysafferentation twice a week and correction made when necessary. Recurrent subluxation patterns are common among babies who react to cow's milk protein, presumably owing to activation of the viscerosomatic reflex by the offending protein. If the neurological dysafferentation is left uncorrected, the child's symptoms may persist, confusing the clinical outcome of the dairy-free trial period. If the child is clinically improved after the 15-day trial period and the correction of the neurological dysafferentation is stable, the mother should be counseled in relation to adopting a dairy-free diet with adequate supplementation for the duration of her breastfeeding, since human milk offers the best protection for the cow's milk reactive infant (Arato et al 1996, Gruskay 1982)
- Artificially fed infants are somewhat more difficult to manp0425 age. Most artificially fed infants who have cow's milk intolerance or allergy will require feeding with a fully hydrolyzed formula, making a 14-21 day trial on such a formula the ideal approach to diagnostic confirmation. However, in some jurisdictions, Government health authorities require an initial trial on a soy-based formula prior to the trial on a fully hydrolyzed formula in order to permit the writing of a formal prescription for the hydrolyzed formula. A significant percentage of children who are allergic or intolerant to cow's milk protein can tolerate soy protein without developing atopic symptoms in the short term (Nadasdi 1992). There are, however, a percentage of children who have a cross-reactivity to soy in whom symptomatic relief will be apparent within the 14-21 day trial, only to relapse again within a few days. Relief comes as the offending cow's milk proteins are eliminated from the diet and exacerbation occurs as a new antigen-antibody response specific to the soy protein begins. It needs to be pointed out that prolonged use of soy infant formula is not a perfect solution; the known problems with such an approach are more fully discussed in Chapter 19.
- p0430 If soy cross reactivity is identified, the child should then be trialed on a hydrolyzed formula such as Nutramigen or Pregestimil. Hydrolyzed formulae consist of dried glucose syrup, casein hydrolysate, a variety of vegetable oils, and a range of

vitamins, minerals and amino acids. Cross-reactivity of hydrolyzed formulae with cow's milk protein has been demonstrated (Ragano et al 1993) but the incidence is very low (Host & Samuelsson 1988, Rugo et al 1992). The major problem encountered with hydrolyzed formulae is rejection by the infant. Cross-reactive infants generally have a high incidence of chronic subluxation recurrence and therefore need to be assessed on a very regular basis during the period of dietary manipulation.

As an aside, casein hydrolysate formulae are much to be p0435 preferred for their hypoallergenic properties than formulae with whey hydrolysate, which demonstrates a greater cross-reactivity with cow's milk (Host & Samuelsson 1988, Rugo et al 1992).

Children who have been shown to be cow's milk allergic or p0440 intolerant should ideally be kept permanently on a dairy free diet. While this strategy has not been demonstrated to completely prevent later development of atopic disease (Bishop et al 1990), there is persuasive evidence suggesting that dietary avoidance of cow's milk protein lessens the incidence of such problems (Schrander et al 1992, Strobel 1992).

Carbohydrate intolerance

s0060

Gastrointestinal intolerance to sugars is thought to affect most p0445 of the adults in the world (Bayless et al 1975) to a greater or lesser extent and is also commonly encountered in infants and older children.

Pathophysiology

s0065

s0070

The principal carbohydrate in human milk is the disaccharide p0450 lactose, which is hydrolyzed in the small intestine by lactase phlorizin hydrolase, otherwise known as lactase (Buller et al 1991). The absence of lactase permits the passage of undigested lactose into the large intestine and is associated with the well-known syndrome referred to as lactose intolerance.

Clinical presentation

The most common presentation of lactose intolerance in p0455 infancy almost invariably follows a bout of acute gastroenteritis in which the gastrointestinal mucosa has become damaged. This condition is self-limiting and will improve over time, but it is worth treating in order to gain symptomatic relief.

In older children the symptoms of lactose intolerance usup0460 ally develop gradually beginning several years after birth and presenting with abdominal cramps, bloating, chronic diarrhea, and excessive passage of flatus related to the ingestion of dairy products (Mitchell et al 1975).

While lactose is by far the most common carbohydrate to p0465 cause the above symptom pattern, some children will experience similar symptoms of abdominal distress and severe diarrhea due to sucrase isomaltase deficiency. This condition is very uncommon and need only be considered in the event that management implemented for lactose intolerance fails.

The irritable baby

CHAPTER 5

Diagnosis

p0470 The diagnosis of lactose intolerance is a clinical one dependent on the recognition of the symptom complex described above and does not normally require any laboratory investigations. However, simple tests which can be carried out on the liquid portion of the diarrheal stool may be performed if there is doubt about the diagnosis. Immediately after collection, the liquid stool specimen is mixed with two parts water, and 15 ml of the resultant fluid is tested with two Clinitest tablets for the presence of reducing sugars, while another drop is applied to the glucose tester on a Clinistix strip. Glucose at 0.5% or less should be considered normal (Behrman et al 2007). In addition, it is worthwhile to test the pH, which will be 5.5 or less in lactose-intolerant individuals.

Management

s0085

- p0475 In breastfed infants, neurological dysafferentation should be corrected and the mother strongly encouraged to continue nursing. Weaning to a lactose-free formula should only be considered in the most severe, protracted cases where there is demonstrable weight loss. In artificially fed infants, the formula should be changed to one that is lactose free.
- p0480 Older children in whom lactose intolerance has made a gradual appearance should be placed on a dairy-free diet, provided with adequate nutritional supplementation, and have their patterns of neurological dysafferentation corrected and monitored at regular intervals. Symptomatic resolution will come quickly and permanently provided there are no 'binges' on dairy products in which as little as 8 ounces of milk (half a glass) may produce a flat blood sugar curve, bloating, cramps, loose stools, or diarrhea (Mitchell et al 1975).

Gastroesophageal reflux

- p0485 Gastroesophageal reflux (GER) is a common occurrence during infancy with most children having a small degree of reflux that is of little clinical consequence, usually requiring only minimal intervention in the event it causes harm (Catto-Smith 1998, Glassman et al 1995). This phenomenon is due to the fact that the intra-abdominal segment of the esophagus is virtually nonexistent at birth (Hull & Johnston 1987) and therefore no effective reflux barrier exists. This barrier is created over the first few months of life as the intra-abdominal esophageal segment lengthens. It is important to recognize that the natural history of GER in infants differs significantly from reflux in adults (Catto-Smith 1998). In children, refluxinduced injury can result from acid exposure, nutrient loss, or respiratory complications.
- p0490 GER has become a rather 'fashionable' diagnosis in recent years in infants that are irritable, with the emergence of the totally unfounded notion of 'silent reflux' in which it is hypothesized that acid reflux into the distal esophagus occurs without the visible evidence of frequent and excessive small volume vomitus. This notion lacks any scientific evidence at all.

Clinical presentation

In true GER the gastrointestinal signs and symptoms are p0495 related directly to the exposure of the distal esophageal epithelium to refluxed gastric contents (Behrman et al 2007). The majority of children with GER have delayed gastric emptying, and vomiting may be forceful because of pylorospasm (Behrman et al 2007). In the vast majority of children, excessive vomiting occurs within the first week of life (Behrman et al 2007, Smith 1980). Vomiting due to GER can occur at any time, unlike normal posseting in which infants will bring up a little milk as they expel swallowed air after feeding.

Respiratory systems are common in children with GER, p0500 usually owing to aspiration of refluxed material. Symptoms include chronic cough, wheezing, and recurrent pneumonia. To be safe, GER should always be considered as a likely etiology of recurrent respiratory symptoms in the first 2 years of life and appropriate diagnostic evaluation carried out.

In children with GER, growth and weight gain are usually p0505 adversely affected. This has been demonstrated in about twothirds of all cases with confirmed GER (Behrman et al 2007).

Diagnosis

In mild cases, diagnosis is made on history and clinical assess- p0510 ment alone. Episodic vomiting beginning in the first 1–2 weeks of life should underpin a diagnosis of mild GER in infancy. Careful monitoring of the child's response to management will quickly confirm the diagnosis. In more severe cases, it is wise to refer the child to a pediatric specialist for esophagoscopy or barium esophagography under fluoroscopic control. Strictures, recurrent reflux and ragged mucosal outline suggestive of esophagitis are readily seen with barium esophagography; however, esophagoscopy with biopsy is the preferred technique for demonstrating esophagitis.

Management

In mild cases the infant should be neurologically managed and p0515 monitored for symptomatic change over a period of 2 weeks. In the event that the cause of the reflux symptoms is functional the child will show rapid improvement related directly to the sustainability of the correction of patterns of neurological dysafferentation. In cases in which improvement is not sustained and the dysafferentation patterns are recurrent, concurrent treatment with an antacid that does not contain aluminum hydroxide is recommended along with the formula thickening in bottlefed infants.

In more intractable cases, it is appropriate to refer the p0520 affected child to the family general practitioner for the prescription of one of the upper gastrointestinal motility enhancing agents which are generally available around the world today.

In cases involving retarded growth factors, respiratory symp- p0525 toms or excessive intractable vomiting which fails to respond to conservative care, referral to a pediatric specialist should be made in order to arrange for appropriate clinical investigation and possible surgical intervention (Smith 1980).

s0095

s0090

CHAPTER 5

The irritable baby

sound Infantile hypertrophic pyloric stenosis

- p0530 Infantile hypertrophic pyloric stenosis (IHPS) is the most common condition affecting infants that requires surgical intervention (St Peter & Ostlie 2008). The etiology of IHPS is yet to be fully elicited. Since the 1990s, a sharp decline in IHPS has been reported in various countries. A possible correlation has been suggested between the 'Back to Sleep' campaign and the falling incidence of IHPS as its decline parallels that of sudden infant death syndrome (SIDS). Recent research from Scotland, however, has shown that the beginning of the decline in the incidence of IHPS predated by some 2 years that of SIDS (which did follow the 'Back to Sleep' campaign), making a direct correlation unlikely (Sommerfield et al 2008).
- Infants with pyloric stenosis generally present with vomitp0535 ing beginning as early as the end of the first week of life and as late as the fifth month (Behrman et al 2007) with the usual time of onset being around the third week. The vomiting may not, at first, be projectile but will progress to that in time. The development of pyloric stenosis is a progressive and dynamic process. The rate of hypertrophy to the point of meeting diagnostic criteria is unknown and there are no data published in the literature regarding the role of repeat ultrasound in patients with persistent symptoms (Keckler et al 2008). The usual pattern of vomiting is a large volume, forceful vomit following the appearance of an obvious peristaltic wave moving across the upper abdomen. The peristaltic wave is often referred to as 'golf ball peristalsis' as it has the appearance of a moving golf ball located within the abdomen. This wave phenomenon is best visualized by looking tangentially across the abdomen. An olive-shaped mass is also usually palpable midway between the umbilicus and the costal margin, just inferior to the liver border. This mass will normally be more readily palpable immediately following an episode of vomiting.
- IHPS has been linked to inappropriate neuromuscular p0540 action, poor breastfeeding performance, and maternal stress (Behrman et al 2007). The striking success that some children receive from chiropractic bears testimony to an etiology of subluxation complex-induced neuromuscular incoordination. The gastrointestinal tract receives sympathetic supply from T8-L1 and, with the exception of the distal colon, parasympathetic supply via the Xth cranial nerve. Neurological dysafferentation involving the upper cervical complex may affect the function of the Xth cranial nerve, possibly owing to the anatomical proximity of the inferior vagal ganglion (Lawrence 1991), the effect of increased dural tension and altered cerebrospinal fluid pressure gradients. The combination of these factors may potentially affect the function of the gastrointestinal tract. In addition, it has been reported by Wiles (1990) that palpatory evidence of reduced motion in the vertebral segments between T7 and T12 has been found in association with generalized gastrointestinal disease. The diagnosis is made clinically and there is usually no need for the diagnostic imaging procedures to be preformed.
- p0545 Care must be taken when making management decisions in relation to children with pyloric stenosis. As the vomiting continues, a progressive loss of fluid, hydrogen, and chloride ions

occurs which will, if not corrected, lead to hypochloremic metabolic alkalosis. Serum potassium levels usually remain unaffected, but in severe cases a total body potassium deficit may occur. Where no evidence of dehydration or physiological stress from biochemical deficiency can be demonstrated, chiropractic management can be safely instigated. The care program should include careful neurological assessment with correction of any dysafferentation and maternal counseling in relation to feeding procedures and stress reduction. If the likelihood is that the mother is probably not going to be able to significantly change her circumstances to reduce stress levels, a referral after subluxation correction to a hospital for both mother and child is appropriate. In the hospital environment, the mother will receive the help she needs to cope with the demands of her baby and may be able to get some muchneeded rest if sleep deprivation has been an issue. Stabilization of the neurological dysafferentation may ultimately depend on correcting breastfeeding errors and reducing immediate stress levels (Homewood 1981).

In the event that there is little or no improvement in the p0550 very short term after neurological correction as been achieved, the child should be referred to a pediatrician for investigation and possible surgical intervention. IHPS, which is related to neuromuscular incoordination, generally responds quickly and permanently to neurological correction. Failure to do so clearly implies that there is another cause of the neuromuscular incoordination unrelated to the subluxation.

When it is clear that a child is demonstrating evidence p0555 of dehydration or physiological stress arising from biochemical deficiency, a pediatric specialist referral is essential. This state does not, however, contraindicate the delivery of chiropractic care, which should be carried out prior to the referral being effected. The usual medical treatment is preoperative correction of the fluid, acid–base and electrolyte losses followed by an operative approach utilizing the application of laparoscopic or circumumbilical techniques to facilitate the pyloromyotomy (Aldridge et al 2007, Ostlie et al 2004, St Peter et al 2006, St Peter & Ostlie 2008).

Intussusception

Intussusception is defined as 'invagination of bowel into an p0560 adjacent lower segment' (Hull & Johnston 1987). While not a particularly common condition in terms of incidence, it does represent the most frequent cause of bowel obstruction in the first 2 years of life. The typical symptom pattern is characterized by episodic, uncontrollable crying, inactivity, irritability and an unwillingness to be handled, sudden onset of pallor which becomes persistent, vomiting, and occasionally rectal bleeding which may be identified in digital examination of the rectum.

s0110

When a child presents which this confluence of symptoms p0565 along with a tender palpable mass in the abdomen which is usually located in the right upper quadrant or close to the left side of the umbilicus, pediatric referral should be made without performing any investigations. Intussusception is a clinical diagnosis. Children presenting to chiropractors with symptoms and signs consistent with this condition should be referred for

The irritable baby

CHAPTER 5

urgent surgical care as bowel viability can be quickly lost (Hull & Johnston 1987).

p0570 Intussusception in a baby that has not been seen previously may be more readily recognizable than is the case with an irritable baby that is being given ongoing chiropractic care, particularly if the symptomatic response is favorable. The latter may be present in such a way that both mother and chiropractor are led into believing that the baby is simply undergoing another paroxysm of irritability caused by a recurrence of the neurological dysafferentation. The critical decision-making problem in this case is linked to the fact that the viscerosomatic reflex responsible for neurological dysafferentation with 'colic' may well be activated in an identical manner by the intussusception. To avoid making an incorrect clinical decision, the abdomen of the irritable baby should be carefully examined at each review consultation.

References

- Aldridge, R.D., MacKinlay, G.A., Aldridge, R.B., 2007. Choice of incision: the experience and evolution of surgical management of infantile hypertrophic pyloric stenosis. J. Laparoendosc. Adv. Surg. Tech. A 17 (1), 131–136.
- Ali, S., Hayek, R., Holland, R., et al., 2002. Effect of Chiropractic Treatment on the Endocrine and Immune System in Asthmatic Patients, Proceedings of the 2002 International Conference on Spinal Manipulation. Foundation for Chiropractic Education and Research, Des Moines Iowa.
- Anrig, C.A., Plaugher, G., (Eds.) 1998. Pediatric Chiropractic, Williams & Wilkins, Baltimore.
- Arato, A., Szalai, K., Tausz, I., et al., 1996. Favourable effect of breast feeding and late introduction of cow's milk in the prevention of suspected allergic symptoms in infancy. Orv. Hetil. 137 (36), 1979–1982.
- Barau, E., Dupont, C., 1994. Allergy to cow's milk proteins in mothers' milk or in hydrolysed cow's milk infant formulas as assessed by intestinal permeability measurements. Allergy 49 (4), 295–298.
- Barr, R.G., 1998. Crying in the first year of life: good news in the midst of distress. Child Care Health Dev. 24 (5), 425–39.
- Bayless, T.M., Rothfield, B., Massa, C., et al., 1975. Lactose and milk intolerance: clinical implications. N. Engl. J. Med. 292 (22), 1156–1159.
- Behrman, R.E., Jenson, H.B., Nelson, W.E., et al., 2007. Nelson Textbook of Pediatrics, eighteenth ed. WB Saunders, Philadelphia.
- Biedermann, H., 1992. Kinematic imbalance due to suboccipital strain in newborns. J. Man. Med. 6, 151–156.
- Bishop, A.M., Hill, D.J., Hosking, C.S., 1990. Natural history of cow milk allergy: clinical outcome. J. Pediatr. 116 (6), 862–867.
- Bjorksten, B., Ahlstedt, S., Bjorksten, F., et al., 1983. Immunoglobulin E and immunoglobulin G4 antibodies to cow's milk in children with cow's milk allergy. Allergy 38 (2), 119–124.
- Bronfort, G., Evans, R.L., Kubic, P., et al., 2002. Chronic pediatric asthma and chiropractic spinal manipulation: a prospective clinical series and randomized clinical pilot study. J. Manipulative Physiol. Ther. 24 (6), 369–377.
- Budgell, B., 1998. A Neurophysiological Rationale for the Chiropractic Management of Visceral Disorders, Seminar Proceedings of the International College of Chiropractic Friday Forum Series, February 6, Sydney, Australia.

- Buller, H.A., Ringe, E.H., Montgomery, R.K., et al., 1991. Clinical aspects of lactose intolerance in children and adults. Scand. J. Gastroenterol. Suppl. 188, 73–80.
- Burr, M.L., Limb, E.S., Maguire, M.J., et al., 1993. Infant feeding, wheezing, and allergy: a prospective study. Arch. Dis. Child. 68 (6), 724–728.
- Businco, L., Benincori, N., Cantani, A., et al., 1985. Chronic diarrhea due to cow's milk allergy: a 4- to 10-year follow-up study. Ann. Allergy 55 (6), 844–847.
- Catto-Smith, A.G., 1998. Gastroesophageal reflux in children. Aust. Fam. Physician 27 (6), 465–469, 472–473.
- Chandra, R.K., 1997. Five year follow up of highrisk infants with family history of allergy who were exclusively breast-fed or fed partial whey hydrosylate, soy, and conventional cow's milk formulas. J. Pediatr. Gastroenterol. Nutr. 24 (4), 380–388.
- Chandra, R.K., Hamed, A., 1991. Cumulative incidence of atopic disorders in high risk infants fed whey hydrosylate, soy, and conventional cow milk formulas. Ann. Allergy 67 (2/1), 129–132.
- Chandra, R.K., Singh, G., Shridhara, B., 1989. Effect of feeding whey hydrosylate, soy and conventional cow milk formulas on incidence of atopic disease in high risk infants. Ann. Allergy 63 (2), 102–106.
- Davies, N.J., 2000. Chiropractic Pediatrics A Clinical Handbook, Churchill Livingstone, Edinburgh,
- Davies, N.J., Jamison, J.R., 2007. Chiropractic management of the irritable baby syndrome. Chiropractic Journal of Australia 37, 25–29.
- deJong, M.H., Scharp-van der Linden, V.T., Aalberse, R.C., et al., 1998. Randomised controlled trial of brief neonatal exposure to cow's milk on the development of atopy. Arch. Dis. Child. 79 (2), 126–130.
- Fallon, J., Edelman, M.G., 1998. Chiropractic care of 401 children with otitis media: a pilot study. Altern. Ther. Health. Med. 4 (2), 93.
- Foucard, T., 1985. Development of food allergies with special reference to cow's milk allergy. Pediatrics 75 (1/2), 177–181.
- Froehle, R.M., 1996. Ear infection: a retrospective study examining improvement from chiropractic care and analyzing for influencing factors. J. Manipulative Physiol. Ther. 19 (3), 169–177.
- Gemmell, H.A., Jacobson, B.H., 1989. Chiropractic manipulation of enuresis: time series descriptive design. J. Manipulative Physiol. Ther. 12 (5), 386–389.

- Glassman, M., George, D., Grill, B., 1995. Gastroesophageal reflux in children: clinical manifestations, diagnosis, and therapy. Gastroenterol. Clin. North. Am. 21 (1), 71–98.
- Goldborough, J., Francis, D., 1983. Dietary management. In: Proceedings of the Second Fisons Food Allergy Workshop. Medicine Publishing Foundation, Oxford, pp. 89–94.
- Gruskay, F.L., 1982. Comparison of breast, cow and soy feedings in the prevention of onset of allergic disease: a 15 year prospective study. Clin. Pediatr. (Phila) 21 (8), 486–491.
- Gustafsson, D., Lowhagen, T., Andersson, K., 1992. Risk of developing atopic disease after early feeding with cow's milk based formula. Arch. Dis. Child. 67 (8), 1008–1010.
- Gutmann, G., 1987. Blocked atlantal nerve syndromes in infants and small children. J. Man. Med. 25, 5–10.
- Hewson, P., Oberklaid, F., Menahem, S., 1987. Infantile colic, distress, and crying. Clin. Pediatr. (Phila) 26 (2), 69–76.
- Hiscock, H., Jordan, B., 2004. Problem crying in infancy. Med. J. Aust. 181 (9), 507–512.
- Homewood, A.E., 1981. The Neurodynamics of the Vertebral Subluxation, Valkyrie Press, St Petersburg, Florida.
- Host, A., Samuelsson, E.G., 1988. Allergic reactions to raw, pasteurized, and homogenized/pasteurized cow milk: a comparison. A double-blind placebocontrolled study in milk allergic children. Allergy 43 (2), 113–118.
- Host, A., Husby, S., Osterballe, O., 1988. A prospective study of cow's milk allergy in exclusively breastfed infants: incidence, pathogenetic role of early inadvertent exposure to cow's milk formula, and characterisation of bovine milk protein in human milk. Acta Paediatr. Scand. 77 (5), 663–670.
- Hull, D., Johnston, D., 1987. Essential Paediatrics, second ed. Churchill Livingstone, Edinburgh.
- Iacono, G., Cavataio, F., Montalto, G., et al., 1998. Intolerance of cow's milk and chronic constipation in children. N. Engl. J. Med. 339 (16), 1100–1104.
- Illingworth, R.S., 1985. Infantile colic revisited. Arch. Dis. Child. 60, 981–985.
- Kahn A Francois, G., Scottiaux, M., et al., 1988. Sleep characteristics in milk-intolerant infants. Sleep 11 (3), 291–297.
- Kahn, A., Mozin, M.J., Rebuffat, E., et al., 1989. Milk intolerance in children with persistent

CHAPTER 5

The irritable baby

sleeplessness: a prospective double-blind crossover evaluation. Pediatrics 84 (4), 595–603.

- Keckler, S.J., Ostlie, D.J., Holcomb III, G.W., et al., 2008. The progressive development of pyloric stenosis: a role for repeat ultrasound. Eur. J. Pediatr. Surg. 18 (3), 168–170.
- Kloughart, N., Nilsson, N., Jacobsen, J., 1989. Infantile colic treated by chiropractors: a prospective study of 316 cases. J. Manipulative Physiol. Ther. 12, 281–288.
- Larson, K., Ayllon, T., 1990. The effects of contingent music and differential reinforcement on infantile colic. Behav Res Ther 28 (2), 119–125.
- Lawrence, D., 1991. Fundamentals of Chiropractic Diagnosis and Management, Williams & Wilkins, Baltimore.
- Lebouef, C., Brown, P., Herman, A., et al., 1991. Chiropractic care for children with nocturnal enuresis: a prospective outcome study. J. Manipulative Physiol. Ther. 14 (2), 110–115.
- Leung, A.K., Lemay, J.F., 2004. Infantile colic: a review. J. R. Soc. Health 124 (4), 162–166.
- Lucas, A., Brooke, O.G., Morley, R., et al., 1990. Early diet of preterm infants and development of allergic or atopic disease: randomised prospective study. Br. Med. J. 300 (6728), 837–840.
- McCarty, E., Frick, O., 1983. Food sensitivity: keys to diagnosis. J. Pediatr. 102 (5), 645–646.
- Meadow, S.R., Smithells, R.W., 1975. Lecture Notes on Pediatrics, second ed. Blackwell Scientific Publications, Oxford.
- Mercer, C., Nook, B.C., 1999. The efficacy of chiropractic spinal adjustments as a treatment protocol in the management of infantile colic, Proceedings of the 5th Biennial Congress, Auckland, New Zealand, May 17–22, pp 170–171.
- Mitchell, K.J., Bayless, T.M., Paige, D.M., et al., 1975. Intolerance of eight ounces of milk in healthy lactose-intolerant teenagers. Pediatrics 56 (5), 718–721.

- Nadasdi, M., 1992. Tolerance of a soy formula by infants and children. Clin. Ther. 14 (2), 236–241. (Ed.), Modern Developments in the
- Nilsson, N., 1985. Infantile colic and chiropractic. Eur. J. Chiropractic 33, 624–625.
- Nilsson, N., Christiansen, B., 1988. Prognostic factors in bronchial asthma in chiropractic practice. J. Aust. Chiropractic Assoc. 18, 85–87.
- Olafsdottir, E., Forshei, S., Fluge, G., et al., 2001. Randomised controlled trial of infantile colic treated with chiropractic spinal manipulation. Arch. Dis. Child. 84 (2), 138–141.
- Ostlie, D.J., Woodall, C.E., Wade, K.R., et al., 2004. An effective pyloromyotomy length in infants undergoing laparoscopic pyloromyotomy. Surgery 136 (4), 827–832.
- Plebani, A., Albertini, A., Scotta, S., et al., 1990. IgE antibodies to hydrosylates of cow's milk in children with cow milk allergy. Ann. Allergy 64 (3), 279–280.
- Ragano, V., Giampietro, P.G., Bruno, G., et al., 1993. Allergenicity of milk protein hydrolysate formulae in children with cow's milk allergy. Eur. J. Pediatr. 152 (9), 760–762.
- Reed, W.R., Beavers, S., Reddy, S.K., et al., 1994. Chiropractic manipulation of primary enuresis. J. Manipulative Physiol. Ther. 17 (9), 596–600.
- Rugo, E., Wahl, R., Wahn, U., 1992. How allergenic are hypoallergenic infant formulae? Clin. Exp. Allergy 22 (6), 635–639.
- St Peter, S.D., Ostlie, D.J., 2008. Pyloric stenosis: from a retrospective analysis to a prospective clinical trial – the impact on surgical outcomes. Curr. Opin. Pediatr. 20 (3), 311–314.
- St Peter, S.D., Holcomb 3rd., G.W., Calkins, C. M., et al., 2006. Open versus laparoscopic pylorotomy for pyloric stenosis: a prospective, randomized trial. Nat. Clin. Pract. Gastroenterol. Hepatol. 4 (4), 196–197.

- Sato, A., 1980. Physiological studies of the somatoautonomic reflexes. In: Haldemann, S. (Ed.), Modern Developments in the Principles and Practice of Chiropractic. Appleton Century Croft, New York.
- Savilahti, E., Kuitunen, M., 1992. Allergenicity of cow milk proteins. J. Pediatr. 121 (5/2), S12–S20.
- Schrander, J.J., Ousden, S., Forget, P.P., et al., 1992. Follow up study of cow's milk protein intolerant infants. Eur. J. Pediatr. 151 (10), 783–785.
- Smith, H.L., 1980. Gastro-oesophageal reflux and hiatal hernia in children. N Z. Med. J. 92 (666), 148–151.
- Sommerfield, T., Chalmers, J., Youngson, G., et al., 2008. The changing epidemiology of infantile hypertrophic pyloric stenosis in Scotland. Arch. Dis. Child. 93 (12), 1003–1004.
- Strobel, S., 1992. Dietary manipulation and induction of tolerance. J. Pediatr. 121 (5/2), S74–S79.
- Taubman, B., 1990. Parental counselling compared with elimination of cow's milk or soy milk protein for the treatment of infantile colic syndrome: a randomized trial. Pediatrics 81 (6), 756–761.
- Wiberg, J.M., Nordsteen, J., Nilsson, N., 1999. The short-term effect of spinal manipulation in the treatment of irritable baby syndrome: a randomized controlled clinical trial with a blinded observer. J. Manipulative Physiol. Ther. 22 (8), 517–522.
- Wiles, M., 1990. Visceral disorders related to the spine. In: Gatterman, M.I. (Ed.), Chiropractic Management of Spine Related Disorders. Williams & Wilkins, Baltimore.
- Wilson, N.W., Hamburger, R.N., 1988. Allergy to cow's milk in the first year of life and its prevention. Ann. Allergy 61 (5), 323–327.
- Wilson, N.W., Self, T.W., Hamburger, R.N., 1990. Severe cow's milk induced colitis in an exclusively breastfed neonate: case report and clinical review of cow's milk allergy. Clin. Pediatr. (Phila) 29 (2), 77–80.